BP-OP-1-6

Initial treatment pattern and prognosis of exocrine pancreatic cancer in Korea from 2006 to 2017

Mee Joo KANG¹, Jiwon LIM², Sung-Sik HAN¹, Hyeong Min PARK¹, Sang-Jae PARK¹, Young-Joo WON^{2,3}, Sun-Whe KIM^{*1}

¹Center for Liver and Pancreatobiliary Cancer, National Cancer Center, Goyang, Korea

²Division of Cancer Registration and Surveillance, National Cancer Center, Goyang, Korea

³Department of Cancer Control & Population Health, Graduate School of Cancer Science and Policy, National Cancer Center, Goyang, Korea

Introduction: Hospital-based database had limitations in holistic analysis of cancer treatment and prognosis, omitting those who are out-of-hospital. In addition, elderly patients with pancreatic cancer were under-represented in clinical studies.

Methods: “First course of treatment (FT)” recorded by cancer registries refers to actually administered treatments within four months of diagnosis. Trends in initial treatment pattern and resultant prognosis of exocrine pancreatic cancer (EPC) in Korea from 2006 to 2017 were investigated based on data of the Korea Central Cancer Registry.

Results: Of all 62,209 EPC cases, localized and regional SEER stage (LR), patients over 70 years-old, and adenocarcinoma and its variants (ADC) accounted for 40.6%, 50.1%, and 95.9%, respectively. The most frequent FT was “no active treatment (NAT)” (46.5%), followed by chemotherapy with/without/or radiotherapy (28.7%) and surgery (22.0%). Among 25,198 LR EPC patients, surgical FT increased from 35.9% to 46.3% and NAT decreased from 45.0% to 29.5%. NAT rate was higher in octogenarian (81.2%) and septuagenarian (44.7%) compared to those younger than 70 years-old (19.6%, $p < 0.001$). Patients with LR ADC had lower rate of surgery (39.6%) compared to cystic cancer variants (86.7%) or acinar cell carcinoma (82.8%, $p < 0.001$). Five-year relative survival rates of LR ADC were higher after surgery (46.1% [localized], 23.6% [regional]) compared to NAT (12.9%, 4.9%, respectively).

Conclusions: Over 55% of patients with LR EPC in their 70s or 80s do not receive active treatment within four months of diagnosis. Clinicians should pay attention to elderly EPC patients to provide proper medical advice.

Corresponding Author: Sun-Whe KIM (sunkim@ncc.re.kr)

Presenter: Mee Joo KANG (meejookang@ncc.re.kr)

Copyright © The Korean Association of Hepato-Biliary-Pancreatic Surgery

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.