EP-16

Perioperative and oncologic outcomes of right anterior sectionectomy for liver disease: A single-center experience with 415 patients

Jae Hoon LEE*, Jaewoo KWON, Yejong PARK, Woohyung LEE, Ki Byung SONG, Dae Wook HWANG, Song Cheol KIM, Young Joo LEE

Division of Hepato-Biliary and Pancreatic Surgery, Department of Surgery, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

Introduction: Right anterior sectionectomy (RAS) is technically difficult and performed infrequently, so there were a few reports of small numbers about this surgery. We described large number of clinicopathologic and oncologic outcomes associated with RAS

Methods: We retrospectively reviewed the medical records of 415 patients treated with RAS for hepatic tumors located at segment 5 and/or 8 between January 2008 and December 2017.

Results: All patients underwent RAS with the alternative Glissonean pedicle clamp and Kelly clamp-crushing methods for transection. The mean operative time was 165 minutes and the mean transection time was 28 minutes. Major morbidities ($\geq$ grade III) occurred in 28 cases (6.7%). Bile leakage occurred in 63 patients (15.1%), but no patients required reoperation. Grade A/B/C post-hepatectomy liver failure occurred in 39/7/0 (9.4%/1.7%/0.0%) patients, respectively. There were no in-hospital mortality caused by postoperative complications. The mean hospital stay was 13.3 days. The most common diagnosis was hepatocellular carcinoma (HCC; $n = 361$, 87.0%), followed by intrahepatic cholangiocarcinoma ($n = 15$, 3.6%), mixed HCC and cholangiocarcinoma ($n = 17$, 4.1%), colorectal cancer liver metastasis ($n = 12$, 2.9%). The mean tumor size was 3.8 cm. Among HCC patients, the 5- and 10- year overall survival (OS) rate was 78.3%, 64.4%, and 5- and 10- year disease-free survival (DFS) rate was 57.2%, 37.7%, respectively. Operative time, tumor size, and vessel invasion were factors significantly associated with OS and DFS for HCC patients.

Conclusions: RAS was associated with acceptable procedure-related morbidity and mortality as well as appropriate oncologic outcomes for HCC patients.

Corresponding Author: Jae Hoon LEE (hbpsurgeon@gmail.com)

Presenter: Jae Hoon LEE (hbpsurgeon@gmail.com)

Copyright © The Korean Association of Hepato-Biliary-Pancreatic Surgery

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.