EP-180

External validation of risk prediction platform for pancreatic fistula after pancreatoduodenectomy using artificial intelligence

So Jeong YOON¹, Jin-Young JANG², So Kyung YOON¹, Ji Hye JUNG¹, Sang Hyun SHIN¹,
Jin Seok HEO¹, Yong Chan SHIN³, Chang-Sup LIM⁴, In Woong HAN^{*1}

¹Division of hepatobiliary-Pancreatic Surgery, Department of Surgery, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

²Department of Surgery, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, Korea

³Department of Surgery, Ilsan Paik Hospital, Inje University College of Medicine, Ilsan, Korea

⁴Department of Surgery, Seoul Metropolitan Government-Seoul National University, Boramae Medical Center, Seoul National University College of Medicine, Seoul, Korea

Introduction: Postoperative pancreatic fistula (POPF) is a life-threatening complication following pancreatoduodenectomy (PD). We previously developed artificial intelligence (AI)-based risk prediction platform for POPF after PD. This study aims to externally validate the formerly developed platform.

Methods: Between January 2007 and December 2016, a total of 1576 patients who underwent PD in Seoul National University Hospital, Ilsan Paik Hospital, and Boramae Medical Center were retrospectively reviewed for external validation. POPF was defined according to the 2016 updated International Study Group of Pancreatic Fistula. The individual risk scores for POPF were calculated using AI risk prediction platform by Samsung Medical Center. The predictive ability was evaluated with a Receiver Operating Characteristic curve and the area under the curve (AUC). In order to obtain the optimal predictive value, we performed backward elimination in accordance with the results from the AI development process.

Results: There were 270 (17.1%) patients who developed POPF in the validation cohort. After backward elimination, each AUC varied from 0.585 to 0.672. The 13 risk factors representing the maximal AUC were pancreatic duct diameter, body mass index, pre-operative serum albumin, amount of intra-operative fluid infusion, age, pre-operative platelet count, location of tumor, combined venous resection, co-existing pancreatitis, pre-operative serum lipase, neoadjuvant radiotherapy, American Society of Anesthesiologists' score, and sex.

Conclusions: We performed external validation of previously developed AI platform for predicting POPF. This user-friendly platform could be used to discriminate individual risk of POPF. Further research is needed to investigate other potential risk factors and to improve the predictability of the platform.

Corresponding Author: In Woong HAN (cardioman76@gmail.com)

Presenter: So Jeong YOON (wooyabi@gmail.com)

Copyright © The Korean Association of Hepato-Biliary-Pancreatic Surgery

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.