EP-6

Liver resection in elderly hepatocellular carcinoma patients (≥ 75 years) is safe

Sunghae PARK, Young Ju OH, Sang Oh YUN, Manuel LIM, Ji Eun JEONG, Eun Sung JEONG,
Jaehun YANG, Jinsoo RHU, Gyu-Seong CHOI, Jong Man KIM*, Jae-Won JOH

Department of Surgery, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

Introduction: As the general population continues to age, elderly patients have dramatically increased for detecting hepatocellular carcinoma (HCC) and required for surgical management. The aim of this study was to compare the postoperative and long-term outcomes of hepatectomy those patients according to the operation time.

Methods: This study included 130 elderly patients who underwent surgical resection for solitary, treatment-naïve HCC between November 1998 and March 2020. Group 1 was defined as patients undergoing LR before 2016 ($n = 68$) and Group 2 was defined as those undergoing liver resection (LR) after 2016 ($n = 62$).

Results: The proportion of major LR and laparoscopic LR in the Group 2 were significantly higher than in the Group 1. Median operation time and blood loss during operation in the Group 2 were smaller than in the Group 1. In addition, intraoperative red blood cell (RBC) transfusion rate, complication rates, and median hospitalization in Group 2 were significantly lower than Group 1. The disease-free survival and patient survival in the Group 2 were better than in the Group 1 even though the presence of tumor grade 3 or 4 and the incidence of microvascular invasion in the Group 1 were higher than in the Group 2. Group 1, long hospitalization, presence of satellite nodule, and intraoperative RBC transfusion were predisposing factors for patient death.

Conclusions: Laparoscopic LR in elderly HCC patients, was performed more frequently than in the past, resulting in an increase in the number of liver surgeries and improved survival.

Corresponding Author: Jong Man KIM (yjongman21@gmail.com)

Presenter: Sunghae PARK (kr6f1207@gmail.com)

Copyright © The Korean Association of Hepato-Biliary-Pancreatic Surgery

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.