LV-OP-2-2

Development and validation of an individualized prediction calculator of postoperative mortality within 6 months after surgical resection for hepatocellular carcinoma: An international multicenter study

Lei LIANG¹, Bing QUAN², Yong-Kang DIAO¹, Chao LI³, Ming-Da WANG³, Wan Yee LAU⁴, Cheng-Wu ZHANG¹, Timothy M. PAWLICK⁵, Dong-Sheng HUANG¹, Feng SHEN³, Tian YANG^{*3}

¹Department of Hepatobiliary, Pancreatic and Minimal Invasive Surgery, Zhejiang Provincial People's Hospital, People's Hospital of Hangzhou Medical College, Hangzhou, China

²Department of Clinical Medicine, Second Military Medical University (Navy Medical University), Shanghai, China

³Department of Hepatobiliary Surgery, Eastern Hepatobiliary Surgery Hospital, Second Military Medical University (Navy Medical University), Shanghai, China

⁴Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China

⁵Department of Surgery, Ohio State University, Wexner Medical Center, Columbus, OH, USA

Introduction: Evidence-based decision-making is critical to optimize the benefits and mitigate futility associated with surgery for patients with malignancies. Untreated hepatocellular carcinoma (HCC) has a median survival of only 6 months. The objective was to develop and validate an individualized patient-specific tool to predict preoperatively the benefit of surgery to provide a survival benefit of at least 6 months following resection.

Methods: Using an international multicenter database, patients who underwent curative-intent liver resection for HCC from 2008 to 2017 were identified. Using random assignment, two-thirds of patients were assigned to a training cohort with the remaining one-third assigned to the validation cohort. Independent predictors of postoperative death within 6 months after surgery for HCC were identified and used to construct a nomogram model with a corresponding online calculator. The predictive accuracy of the calculator was assessed using C-index and calibration curves.

Results: Independent factors associated with death within 6 months of surgery included age, Child-Pugh grading, portal hypertension, alpha-fetoprotein level, tumor rupture, tumor size, tumor number and gross vascular invasion. A nomogram that incorporated these factors demonstrated excellent calibration and good performance in both the training and validation cohorts (C-indexes: 0.802 and 0.798). The nomogram also performed better than four other commonly-used HCC staging systems (C-indexes: 0.800 vs. 0.542–0.748).

Conclusions: An easy-to-use online prediction calculator was able to identify patients at highest risk of death within 6 months of surgery for HCC. The proposed online calculator may help guide surgical decision-making to avoid futile surgery for patients with HCC.

Corresponding Author: Tian YANG (yangtianehbh@smmu.edu.cn)

Presenter: Tian YANG (yangtianehbh@smmu.edu.cn)

Copyright © The Korean Association of Hepato-Biliary-Pancreatic Surgery

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.